

## CircuPlus

The core of **CircuPlus** is a proprietary blend of Green Tea (leaf), Acetyl L-Carnitine, Olive (Leaf), R-Alpha Lipoic Acid, Trans Resveratrol, Coenzyme Q10, and Astaxanthin. It also contains vitamin A (natural mixed carotenoids), vitamin K (phytonadione, and menaquinone-7), vitamin D3 (cholecalciferol), thiamin (benfotiamine), riboflavin (riboflavin 5' phosphate), niacin, vitamin B<sub>6</sub> (P-5-P), folate (L-5 methylfolate calcium), vitamin B<sub>12</sub>, biotin, pantothenic acid, magnesium (magnesium glycerophosphate), zinc (picolinate), selenium (selenomethionine), copper (citrate), and potassium (potassium glycerophosphate). It is designed to ameliorate atherosclerosis and improve circulation.

The core constituents protect against atherosclerosis, improve circulation, and support mitochondrial function. Circulation is supported by Green Tea (leaf)<sup>29, 30, 31,32, 33,34</sup>, Olive (leaf)<sup>22, 23, 24, 25, 26,27, 28</sup>, R-Alpha Lipoic Acid<sup>45, 46, 47, 48, 49</sup>, Astaxanthin<sup>43,44</sup>, and Trans Resveratrol<sup>18, 19, 20, 21</sup>. Mitochondrial function is supported by Coenzyme Q10<sup>66, 67, 68, 69</sup>, Astaxanthin<sup>40, 41, 42</sup>, and Acetyl L-Carnitine<sup>36, 37, 38, 39</sup>.

**CircuPlus** has other constituents that help improve circulation. Vitamin A protects against the oxidation of LDL and slows the progression of atherosclerosis<sup>1, 2</sup>. Menaquinone-7 (vitamin K2) helps prevent calcification of arteries and reduces their stiffness<sup>3, 4, 5, 6, 7</sup>. The product contains vitamins B<sub>6</sub> and B<sub>12</sub>, as well as folate to help lower homocysteine and thereby reduce cardiac risk<sup>8, 50, 51, 52, 53</sup>. Benfotiamine is an S-acyl derivative of thiamine that is more bioavailable than thiamin<sup>15</sup>. Along with being cardioprotective, it helps prevent damage to the arterial endothelium, especially in diabetics<sup>9, 10, 11, 12,13,14, 16, 17</sup>. Vitamin D deficiency can adversely affect the elasticity of arterial walls, so it is included in the product<sup>54, 55</sup>. Niacin reduces inflammation of arterial lining, reduces LDL, and relaxes arterial walls<sup>58,59</sup>. Magnesium lowers cardiovascular risk, reduces endothelial dysfunction, reduces arterial stiffening and reduces cardiovascular risk<sup>60, 61, 62, 63, 64, 65</sup>.

**CircuPlus** is nutritional support for patients with angina, high blood pressure, claudication, erectile dysfunction, and other circulation and cardiac issues.

To place your order for **CircuPlus** or for additional information, please contact us below:

**Zorex International™**

P.O. Box 338 \* Lowell, MI 49331

Toll Free: (800) 679-3027 \* Fax: (800) 257-0319

zorexinternational@gmail.com \* www.zorexinternational.com

CircuPlus

**Recommendations:**

6 capsules daily or as otherwise directed by a health care professional.

**DISCLAIMER:**

Use only as directed. Not for use in pregnancy or lactation. If you have a medical condition, see your doctor or health care provider before using this product. Keep this and all other medications out of the reach of children.

## Supplement Facts

Serving Size: 6 Capsules

Servings Per Container: 30

Amount Per Serving		%DV
Vitamin A (Natural Mixed Carotenoids)	2,100 mcg RAE	233%
Vitamin K (Vit K1 Phytonadione, Vit K2 Menaquinone -7)	50 mcg	42%
Vitamin D3 (Cholecalciferol)	50 mcg	250%
Thiamin (Benfotiamine)	5 mg	417%
Riboflavin (Vit B2)(Riboflavin 5' Phosphate)	5 mg	385%
Niacin (Vit B3)	30 mg	188%
Vitamin B6 (P-5-P)	50 mg	2941%
Folate (L-5-Methylfolate Calcium)	400 mcg DFE	100%
Vitamin B12 (Hydroxycobalamin)	50 mcg	2083%
Biotin	2.5 mg	8333%
Pantothenic Acid (D-Cal Pan)	12.5 mg	250%
Magnesium (Magnesium Glycerophosphate (NutriGP®))	150 mg	36%
Zinc (Zn Picolinate)	15 mg	136%
Selenium (Selenomethionine)	100 mcg	182%
Copper ( Cu Citrate)	500 mcg	56%
Potassium (Potassium Glycerophosphate (NutriGP®))	100 mg	2%
Proprietary Blend	1283 mg	*
Green Tea (Leaf), Acetyl L-Carnitine, Olive (Leaf), R-Alpha Lipoic Acid, Trans Resveratrol, Coenzyme Q10, Astaxanthin		

\* Daily Value (DV) not established

Other Ingredients: NON-GMO Rice Flour, Cellulose, Gelatin.

## References

1. Immunol Invest. 2016;45(1):52-62 **Vitamin A Decreases Cytotoxicity of Oxidized Low-Density Lipoprotein in Patients with Atherosclerosis**
2. J Lipid Res. 2020 Sep 22 (online ahead of print)  **$\beta$ -carotene conversion to vitamin A delays atherosclerosis progression by decreasing hepatic lipid secretion in mice**
3. Nutrients. 2015 Oct 28;7(11):8905-15 **Menaquinone-7 Supplementation to Reduce Vascular Calcification in Patients with Coronary Artery Disease: Rationale and Study Protocol (VitaK-CAC Trial)**
4. Thromb Haemost. 2015 May;113(5):1135-44. **Menaquinone-7 supplementation improves arterial stiffness in healthy postmenopausal women. A double-blind randomised clinical trial**
5. Nutrients. 2015 Aug 18;7(8):6991-7011 **High-Dose Menaquinone-7 Supplementation Reduces Cardiovascular Calcification in a Murine Model of Extrasosseous Calcification**
6. Herz. 2020 Sep;45(6):580-585 **Long-term vitamin-K antagonist use and coronary artery calcification** With its unique selection of patient groups, our study extends the findings of previous research that long-term VKA (vitamin K antagonists) use is related to CAC as detected via CT scanning. The longer and more potent VKA regimen required for MPV patients is the primary cause of CAC (coronary artery calcification) in this group.
7. PLoS One. 2012;7(8):e43229. **Vitamin K-antagonists accelerate atherosclerotic calcification and induce a vulnerable plaque phenotype**
8. J Nutr. 1983 Oct;113(10):2022-33 **Homocystine accumulation in pigs fed diets deficient in vitamin B-6: relationship to atherosclerosis**
9. Mol Cell Biochem. 2009 Jan;320(1-2):149-62. **Ameliorative effect of combination of benfotiamine and fenofibrate in diabetes-induced vascular endothelial dysfunction and nephropathy in the rat**
10. Diabetologia. 2010 Jul;53(7):1506-16 **Increased protein damage in renal glomeruli, retina, nerve, plasma and urine and its prevention by thiamine and benfotiamine therapy in a rat model of diabetes**
11. J Mol Cell Cardiol. 2010 Oct;49(4):625-38 **Benfotiamine improves functional recovery of the infarcted heart via activation of pro-survival G6PD/Akt signaling pathway and modulation of neurohormonal response**
12. Am J Physiol Regul Integr Comp Physiol. 2010 Dec;299(6):R1610-7 **PP2A contributes to endothelial death in high glucose: inhibition by benfotiamine**
13. Circ Heart Fail. 2010 Mar;3(2):294-305 **Vitamin B1 analog benfotiamine prevents diabetes-induced diastolic dysfunction and heart failure through Akt/Pim-1-mediated survival pathway**
14. J Biol Chem. 2006 Apr 7;281(14):9307-13 **Regulation of intracellular glucose and polyol pathway by thiamine and benfotiamine in vascular cells cultured in high glucose**
15. Ann Nutr Metab. 1991;35(5):292-6 **Bioavailability assessment of the lipophilic benfotiamine as compared to a water-soluble thiamin derivative**
16. Ann N Y Acad Sci. 2005 Jun;1043:784-92. **Inhibitors of advanced glycation end product formation and neurovascular dysfunction in experimental diabetes**
17. Sci Aging Knowledge Environ. 2003 Mar 12;2003(10):PE6. **Vitamin B1 blocks damage caused by hyperglycemia**
18. Clin Investig Arterioscler. 2018 Sep-Oct;30(5):209-216. **Effect of resveratrol on seric vascular endothelial growth factor concentrations during atherosclerosis**
19. J Med Food. 2008 Dec;11(4):610-4 **Beneficial effects of resveratrol on atherosclerosis**
20. J Med Food. 2014 Feb;17(2):278-83. **Resveratrol supplementation reduces aortic atherosclerosis and calcification and attenuates loss of aerobic capacity in a mouse model of uremia**
21. Nutr Metab (Lond). 2020 May 27;17:41 **Resveratrol ameliorates atherosclerosis induced by high-fat diet and LPS in ApoE<sup>-/-</sup> mice and inhibits the activation of CD4<sup>+</sup> T cells**
22. Planta Med. 2015 Jun;81(8):648-54 **Effects of the olive tree leaf constituents on myocardial oxidative damage and atherosclerosis**
23. Eur J Nutr. 2008 Aug;47(5):235-43 **The anti-atherosclerotic effect of olive leaf extract is related to suppressed inflammatory response in rabbits with experimental atherosclerosis**
24. Biomed Pharmacother. 2018 Feb;98:446-453 **Olive oil and leaf extract prevent fluoxetine-induced hepatotoxicity by attenuating oxidative stress, inflammation and apoptosis**

25. J Vet Med Sci. 2018 Dec 11;80(12):1801-1807 **Anti-inflammatory effects of olive-derived hydroxytyrosol on lipopolysaccharide-induced inflammation in RAW264.7 cells**
26. Food Chem Toxicol. 2017 Aug;106(Pt A):455-465 **Olive phenolic compounds attenuate deltamethrin-induced liver and kidney toxicity through regulating oxidative stress, inflammation and apoptosis**
27. Animals (Basel). 2019 Jul 7;9(7):427. **Whole Blood Transcriptome Analysis Reveals Positive Effects of Dried Olive Pomace-Supplemented Diet on Inflammation and Cholesterol in Laying Hens**
28. Biomed Pharmacother. 2017 Jul;91:338-349 **Olive (*Olea europaea*) leaf methanolic extract prevents HCl/ethanol-induced gastritis in rats by attenuating inflammation and augmenting antioxidant enzyme activities**
29. J Nutr. 2001 Jan;131(1):27-32 **Tea catechins prevent the development of atherosclerosis in apolipoprotein E-deficient mice**
30. Ann Epidemiol. 2000 Aug;10(6):401-8 **Relation between green tea consumption and the severity of coronary atherosclerosis among Japanese men and women** Green tea protects against atherosclerosis
31. Arterioscler Thromb Vasc Biol. 2010 Oct;30(10):2059-66 **Coffee, decaffeinated coffee, caffeine, and tea consumption in young adulthood and atherosclerosis later in life: the CARDIA study** Tea related to reduced coronary artery plaque
32. Vascul Pharmacol. 2006 Jun;44(6):461-3. **Effect of green tea on angiogenesis and severity of atherosclerosis in cholesterol-fed rabbit**
33. PLoS One. 2017 Aug 4;12(8):e0181666. **Green tea polyphenol treatment attenuates atherosclerosis in high-fat diet-fed apolipoprotein E-knockout mice via alleviating dyslipidemia and up-regulating autophagy**
34. Nutr Res. 2012 Sep;32(9):684-93. **Green tea extract reverses endothelial dysfunction and reduces atherosclerosis progression in homozygous knockout low-density lipoprotein receptor mice**
35. Arch Intern Med. 1999 Oct 11;159(18):2170-4 **Tea flavonoids may protect against atherosclerosis: the Rotterdam Study**
36. Biol Trace Elem Res. 2018 Aug;184(2):422-435. **Acetyl-L-Carnitine Attenuates Arsenic-Induced Oxidative Stress and Hippocampal Mitochondrial Dysfunction**
37. Mech Ageing Dev. 1995 Oct 13;84(2):103-12. **Carnitine-acylcarnitine translocase activity in cardiac mitochondria from aged rats: the effect of acetyl-L-carnitine**
38. Mech Ageing Dev. 2012 Feb-Mar;133(2-3):99-106 **Acetyl-L-carnitine supplementation reverses the age-related decline in carnitine palmitoyltransferase 1 (CPT1) activity in interfibrillar mitochondria without changing the L-carnitine content in the rat heart**
39. FEBS Lett. 1994 Aug 22;350(2-3):213-5 **Effect of aging and acetyl-L-carnitine on the activity of cytochrome oxidase and adenine nucleotide translocase in rat heart mitochondria**
40. Mar Drugs. 2016 Feb 5;14(2):35. **Potential Anti-Atherosclerotic Properties of Astaxanthin** Astaxanthin has been reported to inhibit low-density lipoprotein (LDL) oxidation and to increase high-density lipoprotein (HDL)-cholesterol and adiponectin levels in clinical studies. Accumulating evidence suggests that astaxanthin could exert preventive actions against atherosclerotic cardiovascular disease (CVD) via its potential to improve oxidative stress, inflammation, lipid metabolism, and glucose metabolism.
41. Nutrients. 2018 Feb 28;10(3):281 **Effects of Dietary Supplementation of Astaxanthin and Sesamin on Daily Fatigue: A Randomized, Double-Blind, Placebo-Controlled, Two-Way Crossover Study**
42. J Anim Sci. 2013 Jan;91(1):268-75. **Astaxanthin modulates age-associated mitochondrial dysfunction in healthy dogs**
43. Plant Foods Hum Nutr. 2011 Nov;66(4):363-9 **Positive effects of astaxanthin on lipid profiles and oxidative stress in overweight subjects**
44. Atherosclerosis. 2010 Apr;209(2):520-3 **Administration of natural astaxanthin increases serum HDL-cholesterol and adiponectin in subjects with mild hyperlipidemia**
45. Life Sci. 2010 Jan 16;86(3-4):95-102 **Lipoic acid effects on established atherosclerosis** Magnetic Resonance Imaging (MRI) analysis demonstrated that LA reduced atherosclerotic plaques in the abdominal aorta, with morphological analysis revealing reduced lipid and inflammatory cell content.
46. Lipids Health Dis. 2012 Oct 31;11:148 **Flaxseed oil and  $\alpha$ -lipoic acid combination reduces atherosclerosis risk factors in rats fed a high-fat diet**
47. Diabetes. 2006 Aug;55(8):2238-44  **$\alpha$ -Lipoic acid prevents the increase in atherosclerosis induced by diabetes in apolipoprotein E-deficient mice fed high-fat/low-cholesterol diet**
48. Cell Physiol Biochem. 2018;47(6):2261-2277 **Alpha-Lipoic Acid Protects Human Aortic Endothelial Cells Against H<sub>2</sub>O<sub>2</sub>-Induced Injury and Inhibits Atherosclerosis in Ovariectomized Low Density Lipoprotein Receptor Knock-Out Mice**
49. J Clin Hypertens (Greenwich). 2007 Apr;9(4):249-55 **Effect of combined treatment with alpha-Lipoic acid and acetyl-L-carnitine on vascular function and blood pressure in patients with coronary artery disease**
50. J Nutr Sci Vitaminol (Tokyo). 2019;65(1):1-7 **Relationship between Homocysteine, Folate, Vitamin B 12 and Physical Performance in the Institutionalized Elderly**

51. Neurology. 2013 Oct 8;81(15):1298-307 **Vitamin B supplementation, homocysteine levels, and the risk of cerebrovascular disease: a meta-analysis**
52. Blood Purif. 2006;24(4):379-86 **Homocysteine-lowering vitamin B treatment decreases cardiovascular events in hemodialysis patients**
53. BMC Cardiovasc Disord. 2017 Jan 21;17(1):37 **Serum high concentrations of homocysteine and low levels of folic acid and vitamin B 12 are significantly correlated with the categories of coronary artery diseases**
54. J Cardiovasc Echogr. 2018 Jul-Sep;28(3):171-176. **Impaired Cardiac Functions and Aortic Elastic Properties in Patients with Severe Vitamin D Deficiency**
55. Heart Lung Circ. 2018 Aug;27(8):967-975 **Vitamin D Improves Cardiac Function After Myocardial Infarction Through Modulation of Resident Cardiac Progenitor Cells**
56. Vopr Pitan. 1979 Jan-Feb;(1):26-32 **[Action of biotin-pyridoxine complex on the development of experimental atherosclerosis]**
57. Med Sci Monit. 2015 Dec 29;21:4081-9 **Niacin Suppresses Progression of Atherosclerosis by Inhibiting Vascular Inflammation and Apoptosis of Vascular Smooth Muscle Cells**
58. PLoS One. 2013 Jun 19;8(6):e66467 **Niacin Reduces Atherosclerosis Development in APOE\*3Leiden.CETP Mice Mainly by Reducing Non HDL-Cholesterol**
59. J Cardiovasc Nurs. 2012 Jul-Aug;27(4):303-16. **Niacin-a critical component to the management of atherosclerosis: contemporary management of dyslipidemia to prevent, reduce, or reverse atherosclerotic cardiovascular disease**
60. Int J Mol Sci. 2018 Jun 11;19(6):1724 **Role of Magnesium Deficiency in Promoting Atherosclerosis, Endothelial Dysfunction, and Arterial Stiffening as Risk Factors for Hypertension**
61. J Hypertens. 2017 Jan;35(1):89-97 **Oral magnesium supplementation improves endothelial function and attenuates subclinical atherosclerosis in thiazide-treated hypertensive women**
62. Mol Aspects Med. 2003 Feb-Jun;24(1-3):137-46 **Low magnesium and atherosclerosis: an evidence-based link**
63. Am J Epidemiol. 2009 Jun 15;169(12):1437-44 **Serum and dietary magnesium and risk of ischemic stroke: the Atherosclerosis Risk in Communities Study**
64. Biochim Biophys Acta. 2004 May 24;1689(1):13-21 **Low magnesium promotes endothelial cell dysfunction: implications for atherosclerosis, inflammation and thrombosis**
65. Am J Clin Nutr. 2020 Jan 1;111(1):52-60 **Serum magnesium and the incidence of coronary artery disease over a median 27 years of follow-up in the Atherosclerosis Risk in Communities (ARIC) Study and a meta-analysis**
66. Biochimie. 1999 Dec;81(12):1131-2 **Aging, mitochondria, and coenzyme Q(10): the neglected relationship**
67. FASEB J. 2011 Aug;25(8):2669-87 **Secondary coenzyme Q10 deficiency triggers mitochondria degradation by mitophagy in MELAS fibroblasts**
68. Eur J Pharmacol. 2015 Sep 5;762:270-4 **Coenzyme Q10 remarkably improves the bio-energetic function of rat liver mitochondria treated with statins**